

Specification

Solid selective and differential medium used in isolation and identification of *Salmonella* and coliforms according to the Pharmacopeial Harmonized Method and ISO standard 21150 and 21567.

Formula * in g/L

Pancreatic digest of gelatin	17.000	Sodium chloride	5.000
Meat peptone	1.500	Neutral red	0.030
Casein peptone	1.500	Crystal violet	0.001
Lactose monohydrate	10.000	Agar	15.000
Bile salts	1.500		

Final pH 7.1 ±0.2 at 25 °C

* Adjusted and /or supplemented as required to meet performance criteria

Directions

Suspend 51,5 g of powder in 1 L of distilled water. Bring to the boil and sterilize in the autoclave at 121°C for 15 minutes.

Description

At the beginning of the last century, MacConkey made the original formulation and included ox bile as inhibitor of Gram positive bacteria and litmus as an indicator of acid production from lactose sugar. More recently litmus has been substituted by a phenol red indicator making interpretations easier and more precise. Advancements in the understanding of bacterial physiology has meant that the medium has now been adapted to facilitate the detection of coliforms. The two most significant modifications to the original formulation are as follows:

- The substitution of ox bile by purified bile salts that improves the selectivity and avoids the inherent turbidity, which is due to the fat composition of bile. The efficiency of the inhibition due to bile salts is variable and depends on the relative concentration of cholate and taurocholate.
- The inclusion of supplementary inhibitors such as crystal violet and/or brilliant green. A popular formulation in America, but not in Europe where lower selectivity is preferred.
- Lactose positive bacteria grown on this medium form red colonies due to acid production resulting from lactose fermentation and thus *Escherichia coli* colonies can be easily distinguished as they also form a small precipitation zone of bile salts around them.

Some enterococci may also grow, but they are easy to distinguish from coliforms, as they form smaller colonies and have no precipitation zone.

Technique

Prepare 10-fold serial dilutions of the sample and take 1 ml aliquots from each dilution (in duplicates) and put them into sterile Petri plates. Pour 15 ml of molten medium à 45 °C into every plate and mix carefully. After solidification, a second layer of another 5 ml of sterile medium is poured into every plate to seal the surface and facilitate enumeration of colonies.

For enumeration, after an incubation of 24 hours à 35 °C, select plates with 30-150 colonies. The characteristic colonies must be confirmed as coliforms by gas production from lactose in a broth culture.

Quality control

Incubation temperature: 30-35 °C

Incubation time: 18-72 h

Inoculum: Practical range 100 ±20 CFU. min. 50 CFU (productivity)/ 10⁴-10⁶ CFU (selectivity), according to Ph. Eur. and ISO 11133:2014/Amd 1:2018.

Microorganism

Staphylococcus aureus ATCC® 6538
Enterococcus faecalis ATCC® 29212
Escherichia coli ATCC® 8739
Escherichia coli ATCC® 25922
Salmonella typhimurium ATCC® 14028
Salmonella abony NCTC® 6017
Pseudomonas aeruginosa ATCC® 9027

Growth

Inhibited
 Inhibited
 Productivity > 0.50
 Productivity > 0.50
 Productivity > 0.50
 Productivity > 0.50
 Good

Remarks

Selectivity
 Selectivity
 Dark violet colonies w. precipitate zone
 Dark violet colonies w. precipitate zone
 Colourless colonies without precipitate
 Colourless colonies without precipitate
 Colourless colonies without precipitate

References

- ATLAS, R.M., L.C. PARKS (1993) Handbook of Microbiological Media. CRC Press, Inc. London.
- CLESCERI, L.S., A.E. GEENBERG & A.D. EATON (1998) Standard Methods for the Examination of Water and Wastewater. 20th ed. APHA-AWWA-WEF. Washington. DC. USA.
- DIN 38411-6 (1991) Deutsche Einheitsverfahren zur Wasser-, Abwasser- und Schlammuntersuchung; Mikrobiologische Verfahren (Gruppe K); Nachweis von Escherichia coli und coliformen Keimen (K6)
- DOWNES, F.P. & K. ITO (2001) Compendium of Methods for the Microbiological Examination of Foods. 4th ed. APHA. Washington.
- EUROPEAN PHARMACOPOEIA 11.0 (2023) 11th ed. § 2.6.13. Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. EDQM. Council of Europe. Strasbourg.
- HITCHINS, A.D., P. FENG, W.D. WATKINS, S.R. RIPEY & C.A. CHANDLER (1998) E. coli and coliform bacteria. Bacteriological Analytical Manual. 8th ed. AOAC International. Gaithersburg. MD. USA.
- HORWITZ, W. (2000) Official Methods of Analysis. AOAC Intl. Gaithersburg. MD. USA.
- ISO 11133:2014/ Adm 1:2018. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.
- ISO 21150: 2015 Standard. Cosmetics - Detection of E. coli.
- ISO Standard 21567: 2004. Microbiology of Foods and animal feeding stuffs - Horizontal method for the detection of Shigella spp.
- MacCONKEY, A.T. (1905) Lactose-fermenting Bacteria in faeces. J. Hyg 5:333.
- MURRAY, P.R., E.J. BARON, M.A. PFALLER, F.C. TENOVER, & R.H. YOLKEN (Eds) (1995) Manual of Clinical Microbiology. 6th ed. A.S.M. Washington. DC. USA.
- RAPPAPORT, F. & E. HENING (1952) Media for the isolation and differentiation of pathogenic E. coli (serotypes O111 and O55) J. Clin. Pathology 5:361-362.
- USP 33 - NF 28 (2011) <62> Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. USP Corp. Inc. Rockville. MD. USA.
- VARNAM, A.H. & M.G. EVANS (1991) Foodborne pathogens. Manson Publishing Ltd. London. UK.
- WHO (1963) International Standards for Drinking Waters. 7th ed. Churchill Ltd. London.
- FDA-BAM (2018): Media Index for BAM - BAM Media M91: MacConkey Agar. Food and Drug Administration - Bacteriological Analytical Manual.

Storage

For laboratory use only. Keep tightly closed, away from bright light, in a cool dry place (+4 °C to 30 °C).