

Specification

Sterile selective supplement used for the isolation and enumeration Yeast and Mould from foodstuffs.

Presentation

	Packaging Details	Shelf Life	Storage
10 Freeze dried vials Vial with: 3 ± 0.1 g	23x60 mm glass vials, tag labelled, White plastic cap - 10 vials per box.	49 months	2-25 °C

Composition

Compositon (g/vial)

Oxytetracycline..... 0.050

Reconstitute the original freeze-dried vial
by adding
Sterile distilled water.....6 ml

NOTE : Each vial is sufficient to supplement
500ml of OGYE Agar Base.

Description /Technique

Description:

Oxytetracycline supplement permits to isolate yeasts and moulds in a medium with a neutral pH that gives increased count of them from a variety of foodstuffs compared with media which relied on a low pH to suppress bacterial growth.

In this way physically stressed yeast cells give higher counts on media which depend upon broad-spectrum antibiotics rather than a low pH for selectivity.

Technique:

Collect, dilute and prepare samples and volumes as required according to specifications, directives, official standard regulations and/or expected results.

Reconstitute 1 vial with the sterile diluent (distilled water) in aseptic conditions and add it to 500 ml of OGYE agar base cooled to 50°C temperature.

Do not overheat once supplemented.

Pour the complete medium into Petri dishes and, once solidified on a flat surface, spread the plates by streaking methodology or by spiral method.

Incubate the plates in aerobic atmosphere for 5 days at 20-25 °C, checking for formation of aerial mycelia after 2-5 days.

Incubation times longer than those mentioned above or different incubation temperatures may be required depending on the sample or the specifications.

After incubation, count all the colonies that have appeared onto the surface of the agar.

Presumptive isolation of any pathogenic Yeast and/or Mould must be confirmed by further microbiological and biochemical tests.

Quality control**Physical/Chemical control**

Color : yellow

Microbiological control

Add 1 vial to 500 ml of medium base. DO NOT HEAT once supplemented.

Microbiological control according to ISO 11133:2014/A1:2018.

Analytical methodology according to ISO 11133:2014/A1:2018; A2:2020.

Aerobic. Incubation at 22.5 ± 2 °C until 5 days (moulds and yeast).

Microorganism**Growth**

Aspergillus brasiliensis ATCC® 16404, WDCM 00053

Good

Candida albicans ATCC® 10231, WDCM 00054

Good

S. cerevisiae ATCC® 9763, WDCM 00058

Good

Bacillus subtilis ATCC® 6633, WDCM 00003

Inhibited

Escherichia coli ATCC® 8739, WDCM 00012

Inhibited

Sterility control

100 ml TSB and 100 ml Thioglycollate.

Incubation 48 h at 30-35 °C and 48 h at 20-25 °C: NO GROWTH.

Check at 7 days after incubation in same conditions.

Bibliography

- DOWNES, F.P. & K. ITO (2001) Compendium of methods for the microbiological examination of foods. APHA. Washington DC. USA.
- ISO 11133:2014/ Adm 1:2018. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.
- ISO 13681 Standard (1995) Enumeration of Yeasts and Moulds. Colony Count Technique.
- MacFADDIN, J.F. (1985) Media for isolation-cultivation-identification-maintenance of medical bacteria. William & Wilkins. Baltimore. MD. USA.
- MARSHALL, R.T. (1992) Standard methods for the examination of dairy products 16th ed. APHA. Washington DC, USA.
- MOSSEL, D.A.A., A.M.C. KLEYNEN-SEMMEILING, H.M. VINCENTIE, H. BEERENS & M. CATSARAS (1970) Oxytetracycline-Glucose-Yeast Extract Agar for selective enumeration of moulds and yeasts in foods and clinical material. J. Appl. Bacteriol. 33:454-457.
- SABOURAUD, R. (1910) Les Teignes. Masson, Paris.